



Sharp, G., Stergiakouli, E., Sandy, J., & Relton, C. (2018). Epigenetics and Orofacial Clefts: A Brief Introduction. *Cleft Palate-Craniofacial Journal*, 55(6), 795-797. <https://doi.org/10.1597/16-124>

Publisher's PDF, also known as Version of record

License (if available):
CC BY

Link to published version (if available):
[10.1597/16-124](https://doi.org/10.1597/16-124)

[Link to publication record in Explore Bristol Research](#)
PDF-document

This is the final published version of the article (version of record). It first appeared online via American Cleft Palate-Craniofacial Association at <http://www.cpcjournal.org/doi/abs/10.1597/16-124?code=acpa-premdev#>. Please refer to any applicable terms of use of the publisher.

University of Bristol - Explore Bristol Research

General rights

This document is made available in accordance with publisher policies. Please cite only the published version using the reference above. Full terms of use are available: <http://www.bristol.ac.uk/red/research-policy/pure/user-guides/ebr-terms/>

EDITORIAL

Epigenetics and Orofacial Clefts: A Brief Introduction

Gemma C. Sharp, Ph.D., Evie Stergiakouli, Ph.D., Jonathan Sandy, Ph.D., F.Med.Sci., Caroline Relton, Ph.D.

The term “epigenetics” is used liberally as a possible explanation for traits and conditions that do not appear to be solely genetic in origin. Since orofacial clefts (OFCs) fall into that large category, it is not surprising that the scientific literature in this area has also pointed to a potential etiologic role for epigenetics (Dixon et al., 2011).

WHAT IS EPIGENETICS?

Whereas genetic influences on our traits involve variation in our DNA sequence, epigenetic influences involve changes in how DNA is packaged, expressed, and turned into protein. The underlying DNA remains the same, but chemical modifications change levels of gene activity. These modifications can attach to the DNA itself or to the histone proteins the DNA is wrapped around. In DNA methylation, which is the most commonly studied epigenetic mechanism, methyl groups attach to DNA and may influence expression of nearby genes—often switching them off.

Although every cell in every tissue has essentially the same DNA sequence, epigenetic processes differ between cells. They play a large role in cellular differentiation by enabling a cell to express only the genes that are necessary for its function. Epigenetic modifications can last for the whole of a cell’s life and survive division, but they can also change in response to external influences. Together, this

means that environmental factors might trigger long-term changes in gene activity, which in turn might influence health and disease over an individual’s life (Godfrey et al., 2011). Some evidence also suggests that epigenetic modifications to gamete DNA might enable these traits to be passed to the next generation. Although most of the DNA is stripped of methylation in early development, experiments have shown that marks can remain on certain parts of the DNA that cause aberrant methylation patterns to be reestablished during development (Greer et al., 2014). Another way in which epigenetics can be passed between generations is during pregnancy, when a mother’s environment can influence the epigenome (all of the epigenetic modifications in the genome) of her developing baby *in utero*.

IS EPIGENETICS RELEVANT TO OROFACIAL CLEFTS?

Epigenetic processes, particularly DNA methylation, are important in determining the structure and function of the developing embryonic tissues, so it is plausible that aberrant epigenetic mechanisms may contribute to OFCs. Although human evidence is lacking, evidence from mouse models suggests an important role for DNA methylation and other epigenetic processes in orofacial development and OFCs (Kuriyama et al., 2008; Seelan et al., 2013; Juriloff et al., 2014).

Additionally, in humans, large observational epigenome-wide studies and meta-analyses have found consistent associations between putative risk factors for OFCs and differential methylation in DNA of babies of exposed mothers. For example, maternal obesity (Sharp et al., 2015) and folate deficiency (Joubert et al., 2016a) have been linked to changes in the methylation of neonatal cord blood DNA. Through its role as a coenzyme of one-carbon metabolism, folate is directly involved in methyl group transfer in DNA methylation, which might support evidence for folate deficiency and impaired folate metabolism as risk factors for OFCs. Prenatal tobacco smoke exposure has a particularly large effect on offspring DNA methylation. A recent meta-analysis of epigenome-wide association studies from 13 cohorts (n=6685) identified more than 6000 sites near 3620 genes that were differentially methylated in relation to maternal smoking (Joubert et al., 2016b). Twenty-seven of these genes have previously been linked to OFCs, including *VAX1*, *NOG*, *BMP4*, and *MSX1*. This suggests that maternal smoking may influence offspring risk of OFC through a mechanism involving DNA methylation.

Dr. Sharp is Lecturer in Molecular Epidemiology, MRC Integrative Epidemiology Unit and The Cleft Collective, University of Bristol, Bristol, United Kingdom. Dr. Stergiakouli is Lecturer in Genetic Epidemiology and Statistical Genetics, MRC Integrative Epidemiology Unit and The Cleft Collective, University of Bristol, Bristol, United Kingdom. Dr. Sandy is Dean of Health Sciences, University of Bristol, Bristol, United Kingdom. Dr. Relton is Professor of Epigenetic Epidemiology, MRC Integrative Epidemiology Unit, University of Bristol, Bristol, United Kingdom.

G.C.S., E.S., and C.L.R. are members of the MRC Integrative Epidemiology Unit (IEU) funded by the UK Medical Research Council (MC_UU_12013) and the University of Bristol. G.C.S., E.S., and J.S. are members of the Cleft Collective funded by the Scar Free Foundation (REC approval 13/SW/0064). The views expressed in this publication are those of the author(s) and not necessarily those of The Healing Foundation or The Cleft Collective Cohort Studies team.

This is an open access article distributed under the terms of the Creative Commons Attribution License (CCBY). This license permits copying and redistribution in any medium or format as long as proper attribution is given.

Submitted May 2016; Revised May 2016, June 2016; Accepted July 2016.

Address correspondence to: Dr. Gemma C Sharp, MRC Integrative Epidemiology Unit and The Cleft Collective, University of Bristol, Bristol, BS8 2BN, United Kingdom. E-mail: gemma.sharp@bristol.ac.uk.

DOI: 10.1597/16-124




FIGURE 1 Potential pathways through which epigenetics might influence development of orofacial clefts and cleft-associated outcomes.

Although we are unaware of any studies investigating this, it seems plausible that epigenetic modifications associated with the development of OFCs could also directly influence longer-term outcomes; that is, the downstream consequences of having an OFC may be mediated by changes to the epigenome, and these changes may be independent of the factors that caused the OFC (Fig. 1). Observational studies have found associations between early-life exposures and differential methylation that persists in later life (Simpkin et al., 2015) as well as associations between early-life methylation and later-life OFC-associated outcomes, such as cancer (Verma, 2012).

WHAT ARE THE IMPLICATIONS OF EPIGENETICS FOR CLINICAL PRACTICE?

As technologies to measure epigenetic phenomena become less expensive, and the statistical tools to analyze these become more refined, the potential to use epigenetic analysis in the clinic is increasing. In line with this, two main clinical applications are emerging: (1) development of epigenetic biomarkers to predict or classify disease and (2) exploitation of epigenetic mechanisms to prevent or treat disease.

EPIGENETIC BIOMARKERS FOR PREDICTION OR CLASSIFICATION

Most notably in cancer research, aberrant DNA methylation is being explored as a biomarker of tumor progression for both diagnosis and prognosis (Delpu et al., 2013). One advantage of an epigenetic biomarker over a genetic one is that aberrant methylation is more common than genetic mutation. However, the unique features of epigenetic processes also present challenges in developing accurate biomarkers: (1) some epigenetic changes are cell-specific and may not be detectable in diagnostic tissues such as blood; (2) some epigenetic changes are reversible, so the timing of measurement is important; and (3) epigenetic changes can occur in response to a wide range of environmental factors and are influenced by the underlying genetic sequence, so factors such as age, sex, diet, smoking,

and genotype may affect biomarker efficacy. Nevertheless, epigenetic biomarkers are showing some success as predictive tools, and it is possible that such tools may be useful in predicting or classifying OFCs or downstream consequences of OFCs. For example, if a biomarker can be measured in an accessible tissue, such as amniotic fluid, maternal blood, or chorionic villus blood, it might be a useful tool for antenatal diagnosis or classification of OFCs. Additionally, if measured at birth or early childhood, a biomarker might be useful in predicting surgical success, likelihood of scarring, or longer-term health outcomes, thus identifying groups of patients who might benefit from extra monitoring.

EPIGENETIC MECHANISMS FOR TREATMENT OR PREVENTION

Epigenetic studies in relevant tissue, such as lip/palate tissue collected during cleft surgery, can be combined with causal inference techniques to offer valuable insights into causation of disease and therefore biological mechanisms. Ultimately, this will help to inform intervention and prevention. As mentioned previously, epigenetic changes are reversible and can be influenced by external factors. Thus, disease-causing epigenetic mechanisms could potentially be manipulated to treat or prevent disease. For example, demethylating agents have shown some success in cancer chemotherapy, and there are current efforts to develop drugs that target epigenetic changes in specific genes (Stein, 2014). If epigenetic mechanisms are shown to have a strong causal role in OFCs, then developing prenatal therapies to alter or halt epigenetic changes may be effective in reducing incidence.

SUMMARY

It is biologically plausible that epigenetic changes might be involved in the development of OFCs and in OFC outcomes. Despite this, there has been very little investigation of such changes in humans. This presents exciting and promising opportunities for further research that could inform better prediction, prevention, and management strategies for clinical use.

REFERENCES

- Delpu Y, Cordelier P, Cho WC, Torrisani J. DNA methylation and cancer diagnosis. *Int J Mol Sci.* 2013;14:15029–15058.
- Dixon MJ, Marazita ML, Beaty TH, Murray JC. Cleft lip and palate: understanding genetic and environmental influences. *Nat Rev Genet.* 2011;12:167–178.
- Godfrey KM, Sheppard A, Gluckman PD, Lillycrop KA, Burdge GC, McLean C, Rodford J, Slater-Jefferies JL, Garratt E, Crozier SE, et al. Epigenetic gene promoter methylation at birth is associated with child's later adiposity. *Diabetes.* 2011;60:1528–1534.
- Greer EL, Bees-Sims SE, Brookes E, Spadafora R, Zhu Y, Rothbart SB, Aristizabal-Corrales D, Chen S, Badeaux A, Jin Q, et al. A histone methylation network regulates transgenerational epigenetic memory in *C.elegans*. *Cell Rep.* 2014;10:113–126.
- Joubert BR, den Dekker HT, Felix JF, Bohlin J, Ligthart S, Beckett E, Tiemeier H, van Meurs JB, Uitterlinden AG, Hofman A, et al. Maternal plasma folate impacts differential DNA methylation in an epigenome-wide meta-analysis of newborns. *Nat Commun.* 2016a;7:10577.
- Joubert BR, Felix JF, Yousefi P, Bakulski KM, Just AC, Breton C, Reese SE, Markunas CA, Richmond RC, Xu, C-J, et al. DNA methylation in newborns and maternal smoking in pregnancy: genome-wide consortium meta-analysis. *Am J Hum Genet.* 2016b;98:680–696.
- Juriloff DM, Harris MJ, Mager DL, Gagnier L. Epigenetic mechanism causes Wnt9b deficiency and nonsyndromic cleft lip and palate in the A/WySn mouse strain. *Birth Defects Res A Clin Mol Teratol.* 2014;100:772–788.
- Kuriyama M, Udagawa A, Yoshimoto S, Ichinose M, Sato K, Yamazaki K, Yoshiharu M, Kunio S, Chisato M. DNA methylation changes during cleft palate formation induced by retinoic acid in mice. *Cleft Palate Craniofac J.* 2008;45:545–551.
- Seelan RS, Appana SN, Mukhopadhyay P, Warner DR, Brock GN, Pisano MM, Greene RM. Developmental profiles of the murine palatal methylome. *Birth Defects Res A Clin Mol Teratol.* 2013;97:171–186.
- Sharp GC, Lawlor DA, Richmond RC, Fraser A, Simpkin A, Suderman M, Shihab HA, Lyttleton O, McArdle W, Ring SM, et al. Maternal pre-pregnancy BMI and gestational weight gain, offspring DNA methylation and later offspring adiposity: findings from the Avon Longitudinal Study of Parents and Children. *Int J Epidemiol.* 2015;44:1288–1304.
- Simpkin AJ, Suderman M, Gaunt TR, Lyttleton O, McArdle WL, Ring SM, Tilling K, Davey Smith G, Relton C. Longitudinal analysis of DNA methylation associated with birth weight and gestational age. *Hum Mol Genet.* 2015;24:3752–3763.
- Stein RA. Epigenetic therapies—a new direction in clinical medicine. *Int J Clin Pract.* 2014;68:802–811.
- Verma M. Epigenome-wide association studies (EWAS) in cancer. *Curr Genomics.* 2012;13:308–313.